



Lincolnshire-Prairie View School District 103
Daniel Wright Junior High School

1370 N. Riverwoods Road • Lincolnshire, IL 60069
847/295-1560 • FAX 847/295-7136
dw.d103.org

Scott H. Warren, Ed.D.
Superintendent

Michelle Blackley
Principal

Thomas Herion
Assistant Principal

Melody Littlefair
Assistant Principal

NOTIFICATION OF HEAD LICE

Dear Parents:

A case of head lice has been reported in your child's grade level.

There are three forms of lice- the nit (egg), the nymph, and the adult. Nits are found firmly attached to the hair shaft fairly close to the scalp and are white/yellow in color and oval shaped. They may look like dandruff, but are hard to remove. They take about 1 week to hatch. Nymphs are baby lice and mature into adults in about 7 days. They feed on blood. The adult is about the size of a sesame seed and is grayish-white. They can live up to 30 days and feed on blood.

Incubation period: (the time between exposure to the disease and the appearance of symptoms)
6 to 10 days from laying to hatching of eggs.

Contagious period: (when the disease can be transmitted to another person) As long as the lice are present.

Signs and symptoms: The lice can cause itching usually on the back of the head and around the ears by biting the scalp to feed. The itching is caused by an allergic reaction to the bite of the louse. Your child may have a "tickling feeling" of something moving in the hair, sores on the head caused by scratching, and irritability.

Treatment: Contact your physician, pharmacist, or school nurse for assistance in obtaining a shampoo that is specific for head lice. Follow the instructions carefully. After shampooing with a pediculocide (lice killing) shampoo, you must be sure to remove all the nits with either a fine-toothed comb or with your fingers. If any nits remain, they can hatch and your child can be reinfested.

How this disease is spread: Head lice are spread from one person to another by direct contact with an infested person's head or by contact with lice-infested items, such as hats, hairbrushes, combs, towels, and bedding.

Control of cases: Notify your school and continue checking your students frequently for at least 2 weeks. Head lice must be treated with a medicated shampoo and all the nits must be removed. Wash all personal items such as hats, bedding, towels, combs, etc. in hot water and soap. If the articles cannot be washed, they should be dry-cleaned or placed in a plastic bag for two weeks. Furniture, upholstery, carpeting, mattresses, and pillows should be vacuumed. Anything that cannot be cleaned, like stuffed animals, should be put in a plastic bag and secured for two weeks.

General prevention measures:

Encourage your child to wash his/her hands frequently, advise them not to share hats, combs, brushes, hair ribbons or pillows, to wash their combs and brushes regularly, and to report any itching.

